2002 Advanced Placement Chemistry Exam Section I: Multiple Choice

Time -1 hour and 30 minutes

NO CALCULATORS MAY BE USED WITH SECTION I.

<u>Note</u>: For all questions, assume that the temperature is 298 K, the pressure is 1.00 atmosphere, and solutions are aqueous unless otherwise specified.

Throughout the test the following symbols have the definitions specified unless otherwise noted.

Т	= temperature	т	= molal
Р	= pressure	L, mL	= liter(s), milliliter(s)
V	= volume	g	= gram(s)
S	= entropy	nm	= nanometer(s)
Η	= enthalpy	atm	= atmosphere(s)
G	= free energy	J, kJ	= joule(s), kilojoule(s)
R	= molar gas constant	V	= volt(s)
n	= number of moles	mol	= mole(s)
М	= molar		

Part A

<u>Directions</u>: Each set of lettered choices below refers to the numbered statements immediately following it. Select the one lettered choice that best fits each statement and then fill in the corresponding oval on the answer sheet. A choice may be used once, more than once, or not at all in each set.

Questions 1–2

Consider atoms of the following elements. Assume that the atoms are in the ground state.

- (A) S (B) Ca (C) Ga (D) Sb (E) Br
- 1. The atom that contains exactly two unpaired electrons
- 2. The atom that contains only one electron in the highest occupied energy sublevel

Questions 3-5 refer to the following molecules.

- (A) CO_2 (B) H_2O (C) CH_4 (D) C_2H_4 (E) PH_3
- 3. The molecule with only one double bond
- 4. The molecule with the largest dipole moment
- 5. The molecule that has trigonal pyramidal geometry

Questions 6–7 refer to the following solid compounds.

- (A) $PbSO_4$ (B) CuO (C) $KMnO_4$ (D) KCl (E) $FeCl_3$
- 6. Is purple in aqueous solution 7. Is white and very soluble in water

- (A) Ne (B) Xe (C) O_2 (D) CO (E) NO
- 8. Has an average atomic or molecular speed closest to that of N_2 molecules at 0°C and 1 atm
- 9. Has the greatest density
- 10. Has the greatest rate of effusion through a pinhole

Questions 11-14 refer to the reactions represented below.

- (A) $H_2SeO_4(aq) + 2 Cl^-(aq) + 2 H^+(aq) \rightarrow H_2SeO_3(aq) + Cl_2(g) + H_2O(1)$
- (B) $S_8(s) + 8 O_2(g) \rightarrow 8 SO_2(g)$
- (C) 3 Br₂(aq) + 6 OH⁻(aq) \rightarrow 5 Br⁻(aq) + BrO₃⁻(aq) + 3 H₂O(l)
- (D) $\operatorname{Ca}^{2+}(aq) + \operatorname{SO}_4^{2-}(aq) \rightarrow \operatorname{CaSO}_4(s)$
- (E) $PtCl_4(s) + 2 Cl^{-}(aq) \rightarrow PtCl_6^{2-}(aq)$
- 11. A precipitation reaction
- 12. A reaction that produces a coordination complex
- 13. A reaction in which the same reactant undergoes both oxidation and reduction
- 14. A combustion reaction

Part B

<u>Directions</u>: Each of the questions or incomplete statements below is followed by five suggested answers or completions. Select the one that is best in each case and then fill in the corresponding oval on the answer sheet.

Questions 15-16 relate to the graph below. The graph shows the temperature of a pure substance as it is heated at a constant rate in an open vessel at 1.0 atm pressure. The substance changes from the solid to the liquid to the gas phase.



- 15. The substance is at its normal freezing point at time (A) t_1 (B) t_2 (C) t_3 (D) t_4 (E) t_5
- 16. Which of the following best describes what happens to the substance between t_4 and t_5 ?

- (A) The molecules are leaving the liquid phase.
- (B) The solid and liquid phases coexist in equilibrium.
- (C) The vapor pressure of the substance is decreasing.
- (D) The average intermolecular distance is decreasing.
- (E) The temperature of the substance is increasing.
- 17. In which of the following groups are the three species isoelectronic; i.e., have the same number of electrons?

(C) O²⁻, S²⁻, Cl⁻

(B) Sc, Ti, V^{2+}

- (A) S^{2-}, K^+, Ca^{2+}
- (D) $Mg^{2+}, Ca^{2+}, Sr^{2+}$ (E) Cs, Ba^{2+}, La^{3+}



- 18. The phase diagram for the pure substance X is shown above. The temperature of a sample of pure solid X is slowly raised from 10°C to 100°C at a constant pressure of 0.5 atm. What is the expected behavior of the substance?
 - (A) It first melts to a liquid and then boils at about 70°C.
 - (B) It first melts to a liquid and then boils at about 30°C.
 - (C) It melts to a liquid at a temperature of about 20°C and remains a liquid until the temperature is greater than 100°C.
 - (D) It sublimes to vapor at an equilibrium temperature of about 20° C.
 - (E) It remains a solid until the temperature is greater than 100°C.
- 19. In which of the following species does sulfur have the same oxidation number as it does in H_2SO_4 ?
 - (A) H_2SO_3 (B) $S_2O_3^{2-}$ (C) S^{2-} (D) S_8 (E) SO_2Cl_2
- 20. A flask contains 0.25 mole of $SO_{2(g)}$, 0.50 mole of $CH_{4(g)}$, and 0.50 mole of $O_{2(g)}$. The total pressure of the gases in the flask is 800 mm Hg. What is the partial pressure of the $SO_{2(g)}$ in the flask?

- 21. In the laboratory, $H_{2(g)}$ can be produced by adding which of the following to 1 *M* HCl(*aq*)?
 - $I. 1 M NH_3(aq) \qquad II. Zn(s) \qquad \qquad III. NaHCO_3(s)$
 - (A) I only (B) II only (C) III only (D) I and II only (E) I, II, and III

$$2 \text{ NH}_3 \rightarrow \text{NH}_4^+ + \text{NH}_2^-$$

22. In liquid ammonia, the reaction represented above occurs. In the reaction NH₄⁺ acts as

(A) a catalyst

- (C) the conjugate acid of NH_3
- (E) the oxidizing agent

(D) the reducing agent

$${}^{235}_{92} U + {}^{1}_{0} n \rightarrow {}^{141}_{55} Cs + 3 {}^{1}_{0} n + X$$

- 23. Neutron bombardment of uranium can induce the reaction represented above. Nuclide X is which of the following?
 - (A) ${}^{92}_{35}$ Br (B) ${}^{94}_{35}$ Br (C) ${}^{91}_{37}$ Rb (D) ${}^{92}_{37}$ Rb (E) ${}^{94}_{37}$ Rb
- 24. A compound contains 1.10 mol of K, 0.55 mol of Te, and 1.65 mol of 0. What is the simplest formula of this compound?
 - (A) KTeO (B) KTe_2O (C) K_2TeO_3 (D) K_2TeO_6 (E) K_4TeO_6

 $3 \operatorname{C}_2\operatorname{H}_2(g) \twoheadrightarrow \operatorname{C}_6\operatorname{H}_6(g)$

- 25. What is the standard enthalpy change, ΔH° , for the reaction represented above? (ΔH°_{f} of C₂H_{2(g)} is 230 kJ mol⁻¹; ΔH°_{f} of C₆H_{6(g)} is 83 kJ mol⁻¹.)
 - (A) -607 kJ (B) -147 kJ (C) -19 kJ (D) +19 kJ (E) +773 kJ
- 26. Approximately what mass of CuSO₄•5H₂O (250 g mol⁻¹) is required to prepare 250 mL of 0.10 M copper(II) sulfate solution?
 - (A) 4.0 g (B) 6.2 g (C) 34 g (D) 85 g (E) 140 g

$$2 \operatorname{NO}(g) + \operatorname{O}_2(g) \rightarrow 2 \operatorname{NO}_2(g)$$

- 27. A possible mechanism for the overall reaction represented above is the following.
 - (1) $NO(g) + NO(g) \rightarrow N_2O_2(g)$ slow

(2) $N_2O_2(g) + O_2(g) \rightarrow 2 NO_2(g)$ fast

Which of the following rate expressions agrees best with this possible mechanism?

(A) Rate = $k[NO]^2$ (B) Rate = $k\frac{[NO]^2}{[O_2]}$ (C) Rate = $k\frac{[NO]^2}{[O_2]}$ (D) Rate = $k[NO]^2[O_2]$ (E) Rate = $k[N_2O_2][O_2]$

28. Of the following compounds, which is the most ionic?

- (A) SiCl₄ (B) BrCl (C) PCl₃ (D) Cl_2O (E) CaCl₂
- 29. The best explanation for the fact that diamond is extremely hard is that diamond crystals
 - (A) are made up of atoms that are intrinsically hard because of their electronic structures
 - (B) consist of positive and negative ions that are strongly attracted to each other
 - (C) are giant molecules in which each atom forms strong covalent bonds with all of its neighboring atoms
 - (D) are formed under extreme conditions of temperature and pressure
 - (E) contain orbitals or bands of delocalized electrons that belong not to single atoms but to each crystal as a whole

30. At 25°C aqueous solutions with a pH of 8 have a hydroxide ion concentration, [OH⁻], of (A) $1 \times 10^{-14} M$ (B) $1 \times 10^{-8} M$ (C) $1 \times 10^{-6} M$ (D) 1 M (E) 8 M

$$CS_2(l) + 3 O_2(g) \rightarrow CO_2(g) + 2 SO_2(g)$$

- 31. What volume of $O_{2(g)}$ is required to react with excess $CS_{2(l)}$ to produce 4.0 L of $CO_{2(g)}$? (Assume all gases are measured at 0°C and 1 atm.)
 - (A) 12 L (B) 22.4 L (C) $\frac{1}{3} \times 22.4$ L (D) 2 × 22.4 L (E) 3 × 22.4 L
- 32. Which of the following oxides is a gas at 25°C and 1 atm?
 (A) Rb₂O
 (B) N₂O
 (C) Na₂O₂
 (D) SiO₂
 (E) La₂O₃

Questions 33-34

The graph below shows the titration curve that results when 100. mL of 0.0250 M acetic acid is titrated with 0.100 M NaOH.



mL of 0.100 M NaOH Added

33. Which of the following indicators is the best choice for this titration?

	Indicator	pH Range of <u>Color Change</u>
(A)	Methyl orange	3.2 - 4.4
(B)	Methyl red	4.8 - 6.0
(C)	Bromothymol blue	6.1 - 7.6
(D)	Phenolphthalein	8.2 - 10.0
(E)	Alizarin	11.0 - 12.4

- 34. What part of the curve corresponds to the optimum buffer action for the acetic acid/acetate ion pair?
 - (A) Point V
 - (C) Point Z

- (B) Point X
- (D) Along all of section WY

- (E) Along all of section YZ
- 35. A solution is made by dissolving a nonvolatile solute in a pure solvent. Compared to the pure solvent, the solution
 - (A) has a higher normal boiling point
 - (C) has the same vapor pressure

- (B) has a higher vapor pressure
- (D) has a higher freezing point

(E) is more nearly ideal

2002- page 5

- 36. A sample of a solution of an unknown was treated with dilute hydrochloric acid. The white precipitate formed was filtered and washed with hot water. A few drops of potassium iodide solution were added to the hot water filtrate and a bright yellow precipitate was produced. The white precipitate remaining on the filter paper was readily soluble in ammonia solution. What two ions could have been present in the unknown?
 - (A) $Ag^+(aq)$ and $Hg_2^{2+}(aq)$
 - (C) $Ba^{2+}(aq)$ and $Ag^{+}(aq)$
 - (E) $Ba^{2+}(aq)$ and $Pb^{2+}(aq)$

 $HCO_3^{-}(aq) + OH^{-}(aq) \Leftrightarrow H_2O(l) + CO_3^{2-}(aq)$ $\Delta H = -41.4 \text{ kJ}$

- 37. When the reaction represented by the equation above is at equilibrium at 1 atm and 25°C, the ratio $[CO_3^{2-}]$ $\frac{1}{|\text{HCO}_3^-|}$ can be increased by doing which of the following?

 - (A) Decreasing the temperature

(B) Adding acid

(D) diluting the solution with distilled water

(B) $Ag^+(aq)$ and $Pb^{2+}(aq)$

(D) $Ba^{2+}(aq)$ and $Hg_2^{2+}(aq)$

- (C) Adding a catalyst
- (E) Bubbling neon gas through the solution
- 38. A 0.10 M aqueous solution of sodium sulfate, Na_2SO_4 , is a better conductor of electricity than 0.10 M aqueous solution of sodium chloride, NaCl. Which of the following best explains this observation?
 - (A) Na_2SO_4 is more soluble water than NaCl is.
 - (B) Na_2SO_4 has a higher molar mass than NaCl has.
 - (C) To prepare a given volume of 0.10 M solution, the mass of Na₂SO₄ needed is more than twice the mass of NaCl needed.
 - (D) More moles of ions are present in a given volume of $0.10 M \text{ Na}_2\text{SO}_4$ than in the same volume of 0.10M NaCl.
 - (E) The degree of dissociation of Na_2SO_4 in solution is significantly greater than that of NaCl.



39. On the basis of the solubility curves shown above, the greatest percentage of which compound can be recovered by cooling a saturated solution of that compound from 90°C to 30°C?
(A) NaCl
(B) KNO₃
(C) K₂CrO₄
(D) K₂SO₄
(E) Ce₂(SO₄)₃

- 40. An excess of Mg(s) is added to 100. mL of 0.400 *M* HCl. At 0°C and 1 atm pressure, what volume of H₂ gas can be obtained?
 - (A) 22.4 mL (B) 44.8 mL (C) 224 mL (D) 448 mL (E) 896 mL
- 41. When solid NH₄SCN is mixed with solid Ba(OH)₂ in a closed container, the temperature drops and a gas is produced. Which of the following indicates the correct signs for ΔG , ΔH , and ΔS for the process?

	ΔG	ΔH	ΔS
(A)	—		—
(B)	—	+	—
(C)	—	+	+
(D)	+	_	+
(E)	+	_	—

$H_2(g) + Br_2(g) \Leftrightarrow 2 HBr(g)$

42. At a certain temperature, the value of the equilibrium constant, K, for the reaction represented above is 2.0×10^5 . What is the value of K for the reverse reaction at the same temperature?

(A)
$$-2.0 \times 10-5$$
 (B) $5.0 \times 10-6$ (C) $2.0 \times 10-5$ (D) $5.0 \times 10-5$ (E) $5.0 \times 10-4$

- 43. The atomic mass of copper is 63.55. Given that there are only two naturally occurring isotopes of copper, ⁶³Cu and ⁶⁵Cu, the natural abundance of the ⁶⁵Cu isotope must be approximately
 - (A) 90% (B) 70% (C) 50% (D) 25% (E) 10%

- 44. Which of the following properties generally decreases across the periodic table from sodium to chlorine?
 - (A) First ionization energy
 - (C) Electronegativity

- (B) Atomic mass
- (D) Maximum value of oxidation number

- (E) Atomic radius
- 45. What is the mole fraction of ethanol, C₂H₅OH, in an aqueous solution that is 46 percent ethanol by mass? (The molar mass of C₂H₅OH is 46 g; the molar mass of H₂O is 18 g.)
 - (A) 0.25 (B) 0.46 (C) 0.54 (D) 0.67 (E) 0.75
- 46. The effective nuclear charge experienced by the outermost electron of Na is different than the effective nuclear charge experienced by the outermost electron of Ne. This difference best accounts for which of the following?
 - (A) Na has a greater density at standard conditions than Ne.
 - (B) Na has a lower first ionization energy than Ne.
 - (C) Na has a higher melting point than Ne.
 - (D) Na has a higher neutron-to-proton ratio than Ne.
 - (E) Na has fewer naturally occurring isotopes than Ne.
- 47. Which of the following is a correct statement about reaction order?
 - (A) Reaction order can only be a whole number.
 - (B) Reaction order can be determined only from the coefficients of the balanced equation for the reaction.
 - (C) Reaction order can be determined only by experiment.
 - (D) Reaction order increases with increasing temperature.
 - (E) A second-order reaction must involve at least two different compounds as reactants.
- 48. Sodium chloride is LEAST soluble in which of the following liquids?
 - (A) H_2O (B) CCl_4 (C) HF (D) CH_3OH (E) CH_3COOH

 $\dots\operatorname{Cr}_2\operatorname{O}_7^{2-}(aq)+\dots\operatorname{H}_2\operatorname{S}(g)+\dots\operatorname{H}^+(aq) \twoheadrightarrow \dots\operatorname{Cr}^{3+}(aq)+\dots\operatorname{S}(s)+\dots\operatorname{H}_2\operatorname{O}(l)$

- 49. When the equation above is correctly balanced and all coefficients are reduced to lowest whole-number terms, the coefficient for $H^+(aq)$ is
 - (A) 2 (B) 4 (C) 6 (D) 8 (E) 14
- 50. Which of the following represents acceptable laboratory practice?
 - (A) Placing a hot object on a balance pan
 - (B) Using distilled water for the final rinse of a buret before filling it with standardized solution
 - (C) Adding a weighed quantity of solid acid to a titration flask wet with distilled water
 - (D) Using 10 mL of standard strength phenolphthalein indicator solution for titration of 25 mL of acid solution
 - (E) Diluting a solution in a volumetric flask to its final concentration with hot water

$$3 \text{Cu}(s) + 8 \text{H}^+(aq) + 2 \text{NO}_3(aq) \rightarrow 3 \text{Cu}^{2+}(aq) + 2 \text{NO}(g) + 4 \text{H}_2\text{O}(l)$$

- 51. True statements about the reaction represented above include which of the following?
 - I. Cu(*s*) acts as an oxidizing agent.
 - II. The oxidation state of nitrogen changes from +5 to +2.
 - III. Hydrogen ions are oxidized to form $H_2O(l)$.
 - (A) I only (B) II only (C) III only (D) I and II (E) II and III

52. Propane gas, C_3H_8 , burns in excess oxygen gas. When the equation for this reaction is correctly balanced and all coefficients are reduced to their lowest whole-number terms, the coefficient for O_2 is

- (A) 4 (B) 5 (C) 7 (D) 10 (E) 22
- 53. According to the VSEPR model, the progressive decrease in the bond angles in the series of molecules CH₄, NH₃, and H₂O is best accounted for by the
 - (A) increasing strength of the bonds
 - (B) decreasing size of the central atom
 - (C) increasing electronegativity of the central atom
 - (D) increasing number of unshared pairs of electrons
 - (E) decreasing repulsion between hydrogen atoms
- 54. Which of the following must be true for a reaction for which the activation energy is the same for both the forward and the reverse reactions?
 - (A) A catalyst is present.
 - (B) The reaction order can be obtained directly from the balanced equation.
 - (C) The reaction order is zero.
 - (D) ΔH for the reaction is zero.
 - (E) ΔS for the reaction is zero.

Time(days)	0	1	2	3	4	5	6	7	 10	 20
% Reactant remaining	100	79	63	50	40	31	25	20	10	1

55. A reaction was observed for 20 days and the percentage of the reactant remaining after each day was recorded in the table above. Which of the following best describes the order and the half-life of the reaction?

	Reaction Order	Half-life (days)
(A)	First	3
(B)	First	10
(C)	Second	3
(D)	Second	6
(E)	Second	10

- 56. The boiling points of the elements helium, neon, argon, krypton, and xenon increase in that order. Which of the following statements accounts for this increase?
 - (A) The London (dispersion) forces increase.
 - (B) The hydrogen bonding increases.
 - (C) The dipole-dipole forces increase.
 - (D) The chemical reactivity increases.
 - (E) The number of nearest neighbors increases.

Rate =
$$k[M][N]^2$$

- 57. The rate of a certain chemical reaction between substances M and N obeys the rate law above. The reaction is first studied with [M] and [N] each 1×10^{-3} molar. If a new experiment is conducted with [M] and [N] each 2×10^{-3} molar, the reaction rate will increase by a factor of
 - (A) 2 (B) 4 (C) 6 (D) 8 (E) 16

 $2 \ \mathrm{N_2H_4}(g) + \mathrm{N_2O_4}(g) \rightarrow 3 \ \mathrm{N_2}(g) + 4 \ \mathrm{H_2O}(g)$

- 58. When 8.0 g of N_2H_4 (32 g mol⁻¹) and 92 g of N_2O_4 (92 g mol⁻¹) are mixed together and react according to the equation above, what is the maximum mass of H_2O that can be produced?
 - (A) 9.0 g (B) 18 g (C) 36 g (D) 72 g (E) 144 g
- 59. All of the halogens in their elemental form at 25°C and 1 atm are
 - (A) conductors of electricity(B) diatomic molecules(C) odorless(D) colorless
 - (E) gases

 $2 \operatorname{H_2O}(l) + 4 \operatorname{MnO_4^-}(aq) + 3 \operatorname{ClO_2^-}(aq) \rightarrow 4 \operatorname{MnO_2}(s) + 3 \operatorname{ClO_4^-}(aq) + 4 \operatorname{OH^-}(aq)$

- 60. According to the balanced equation above, how many moles of $\text{ClO}_2^{-}(aq)$ are needed to react completely with 20. mL of 0.20 *M* KMnO₄ solution?
 - (A) 0.0030 mol (B) 0.0053 mol (C) 0.0075 mol (D) 0.013 mol (E) 0.030 mol
- 61. How can 100. mL of sodium hydroxide solution with a pH of 13.00 be converted to a sodium hydroxide solution with a pH of 12.00?
 - (A) By diluting the solution with distilled water to a total volume of 108 mL
 - (B) By diluting the solution with distilled water to a total volume of 200 mL
 - (C) By diluting the solution with distilled water to a total volume of 1.00 L
 - (D) By adding 100. mL of 0.10 M HCl
 - (E) By adding 100. mL of 0.10 M NaOH

62. Which of the following structural formulas represents an isomer of the compound that has the structural formula represented above?



63. Mixtures that would be considered buffers include which of the following?

	I.	0.10 M HCl + 0.10 M N	laCl	
	II.	0.10 <i>M</i> HF + 0.10 <i>M</i> Na	aF	
	III.	0.10 <i>M</i> HBr + 0.10 <i>M</i> N	laBr	
(A) I only	(B) II only	(C) III only	(D) I and II	(E) II and III

64. Ascorbic acid, $H_2C_6H_6O_{6(s)}$, is a diprotic acid with $K_1 = 7.9 \times 10^{-5}$ and $K_2 = 1.6 \times 10^{-12}$. In a 0.005 *M* aqueous solution of ascorbic acid, which of the following species is present in the <u>lowest</u> concentration?

- (A) $H_2O(l)$ (B) $H_3O^+(aq)$ (C) $H_2C_6H_6O_6(aq)$
- (D) $HC_6H_6O_{6}(aq)$ (E) $C_6H_6O_{6}^{2}(aq)$

65. Which of the following substances is LEAST soluble in water?
(A) (NH₄)₂SO₄ (B) KMnO₄ (C) BaCO₃ (D) Zn(NO₃)₂ (E) Na₃PO₄

66. A 2 L container will hold about 4 g of which of the following gases at 0°C and 1 atm?
(A) SO₂
(B) N₂
(C) CO₂
(D) C₄H₈
(E) NH₃

- 67. Which of the following describes the changes in forces of attraction that occur as H₂O changes phase from a liquid to a vapor?
 - (A) H–O bonds break as H–H and O–O bonds form.
 - (B) Hydrogen bonds between H_2O molecules are broken.
 - (C) Covalent bonds between H_2O molecules are broken.
 - (D) Ionic bonds between H⁺ ions and OH⁻ ions are broken.
 - (E) Covalent bonds between H^+ ions and H_2O molecules become more effective.



68. Liquid naphthalene at 95°C was cooled to 30°C as represented in the cooling curve above. From which section of the curve can the melting point of naphthalene be determined?

(A) A (B) B (C) C (D) D (E) E

69. If 200. mL of 0.60 *M* MgCl_{2(aq)} is added to 400. mL of distilled water, what is the concentration of Mg²⁺(aq) in the resulting solution? (Assume volumes are additive.)
(A) 0.20 *M*(B) 0.30 *M*(C) 0.40 *M*(D) 0.60 *M*(E) 1.2 *M*

- 70. Of the following pure substances, which has the highest melting point?(A) S_8 (B) I_2 (C) SiO_2 (D) SO_2 (E) C_6H_6
- 71. In the electroplating of nickel, 0.200 faraday of electrical charge is passed through a solution of NiSO₄. What mass of nickel is deposited?
 - (A) 2.94 g (B) 5.87 g (C) 11.7 g (D) 58.7 g (E) 294 g
- 72. A colorless solution is divided into three samples. The following tests were performed on samples of the solution.

Sample	Test	Observation
1	Add H ⁺ (aq)	No change
2	Add NH ₃ (aq)	No change
3	Add $SO_4^{2-}(aq)$	No change

Which of the following ions could be present in the solution at a concentration of 0.10 M?

(A) $Ni^{2+}(aq)$ (B) $Al^{3+}(aq)$ (C) $Ba^{2+}(aq)$ (D) $Na^{+}(aq)$ (E) $CO_{3}^{2-}(aq)$

$$X(s) \Leftrightarrow X(l)$$

- 73. Which of the following is true for any substance undergoing the process represented above at its normal melting point?
 - (A) $\Delta S < 0$ (B) $\Delta H = 0$ (C) $\Delta H = T\Delta G$ (D) $T\Delta S = 0$ (E) $\Delta H = T\Delta S$

74. A pure, white crystalline solid dissolves in water to yield a basic solution that liberates a gas when excess acid is added to it. On the basis of this information, the solid could be

(A) KNO_3 (B) K_2CO_3 (C) KOH (D) $KHSO_4$ (E) KCl

75. In a saturated solution of $Zn(OH)_2$ at 25°C the value of [OH-] is $2.0 \times 10^{-6} M$. What is the value of the solubility-product constant, K_{sp} , for $Zn(OH)_2$ at 25°C?

(A) 4.0×10^{-18} (B) 8.0×10^{-18} (C) 1.6×10^{-17} (D) 4.0×10^{-12} (E) 2.0×10^{-6}

		% correct by grade				total %	
item	answer	5	4	3	2	1	correct
1	Α	80	71	65	59	46	63
2	C	95	87	77	64	39	70
3	D	91	85	79	72	52	74
4	В	85	70	57	45	29	54
5	Е	90	82	77	72	61	75
6	С	81	67	56	46	35	55
7	D	84	71	62	53	43	61
8	D	81	59	40	26	15	41
9	В	88	80	73	67	55	71
10	Α	91	79	63	49	28	59
11	D	99	98	92	82	49	82
12	Е	69	42	25	16	11	30
13	С	94	84	73	59	40	67
14	В	97	91	82	68	44	74
15	В	90	74	56	42	32	56
16	Α	89	75	60	47	32	58
17	Α	97	91	80	64	35	70
18	D	97	92	79	61	31	69
19	E	97	91	81	65	38	72
20	E	94	85	74	60	36	67
21	В	85	67	51	36	19	48
22	С	8S	73	61	48	32	57
23	D	63	49	38	29	18	37
24	С	98	95	88	79	59	82
25	Α	92	83	73	60	36	66
26	В	89	77	63	44	23	56
27	A	83	63	47	32	16	45
28	E	93	84	74	61	39	68
29	С	86	68	54	39	24	51
30	С	94	87	80	69	42	72
31	A	92	78	60	43	23	56
32	В	89	79	71	64	52	69
33	D	94	86	76	63	43	70
34	A	75	50	32	19	10	34
35	A	91	74	53	39	25	53
36	В	51	35	26	21	17	28
37	A	85	65	44	29	20	45
38	D	97	84	64	47	30	61

item	answer	5	4	3	2	1	correct
39	В	94	87	76	66	53	73
40	D	69	47	25	12	8	29
41	С	8S	63	42	25	16	43
42	В	62	35	20	11	5	24
43	D	78	63	51	40	26	49
44	Е	92	85	76	6S	42	70
45	Α	88	67	44	24	15	44
46	В	8S	74	65	56	38	62
47	С	69	52	39	29	17	39
48	В	85	66	48	35	23	48
49	D	64	39	27	21	16	31
50	С	38	22	14	8	6	16
51	В	88	72	54	34	14	49
52	В	93	88	79	66	39	71
53	D	94	83	67	50	30	62
54	D	79	60	43	29	18	43
55	А	74	63	57	51	38	55
56	А	93	81	66	51	32	62
57	D	92	77	58	37	15	52
58	А	87	72	52	31	12	47
59	В	70	53	40	30	20	40
60	Α	93	81	60	35	15	53
61	С	62	29	13	7	11	22
62	В	57	38	27	18	12	28
63	В	65	37	20	10	8	25
64	E	88	61	34	17	9	38
65	С	81	60	43	33	22	45
66	С	82	61	40	24	17	41
67	В	88	70	52	40	26	52
68	С	89	73	59	46	34	57
69	Α	86	70	47	29	15	46
70	С	S8	38	26	18	15	29
71	В	43	14	6	5	8	13
72	D	45	48	30	21	18	35
73	E	65	43	31	23	20	34
74	В	67	35	21	15	12	27
75	Α	64	30	13	7	7	21

% correct by gradetotal %Average % correct =

VERY EASY (80–100% correct), 2.7% [11, 24]

EASY (60–79% correct), 33.3% [1, 2, 3, 5, 7, 9, 13, 14, 17, 18, 19, 20, 25, 28, 30, 31, 32, 33, 38, 39, 44, 46, 52, 53, 56]

MEDIUM DIFFICULTY (40–59% correct), 40.0% [4, 6, 8, 10, 15, 16, 21, 22, 26, 27, 29, 31, 35, 37, 41, 43, 45, 48, 51, 54, 55, 57, 58, 59, 60, 65, 66, 67, 68, 69]

VERY HARD (0–19% correct), 2.7% [50, 71]

[1/4 of wrong answers subtracted from # correct]

Multiple-Choice		AP Grade							
Score			(%)			(%)			
	1	2	3	4	5				
46–75	0.0	0.0	1.3	15.2	83.5	17.0			
38-45	0.0	0.5	23.3	55.6	20.6	13.2			
26–37	0.2	15.0	64.6	19.0	1.2	24.7			
15–25	14.6	60.4	24.7	0.3	0.0	23.9			
0-14	81.6	22.0	0.6	0.0	0.0	21.2			
Total	20.8	22.0	25.3	14.7	17.2	100			